45th ANNUAL DIAGNOSTIC SLIDE SESSION, 2003 REFERENCES AND DIAGNOSES

MODERATOR: E. Tessa Hedley-Whyte, M.D.

EDITOR: Leroy R. Sharer, M.D.

Case 2004-10

Submitted by: Dr. Juan M. Bilbao, Dr. Tim Feltis and Ms. Beverley Young, University of Toronto, Sunnybrook & Women's College Health Sciences Centre, and the Office of the Chief Coroner for Ontario, Toronto, Ontario, CANADA

Diagnosis: Dissection of the right vertebral artery (V3 segment) and traumatic pseudoaneurysm of the left vertebral artery (V4 segment) distal to the posterior inferior cerebellar artery (PICA).

Comment: The "treatment" mentioned in the protocol was massage given by an aide to a chiropractor, while the chiropractor was out of the office. The dissecting aneurysm of the right vertebral artery extended for 3 to 4 cm, from C3 up to C1. Electron microscopy failed to demonstrate spiraled collagen that is typical for Ehlers-Danlos syndrome in the vessel with the dissection. Dissection has been described previously following neck manipulation, but pseudoaneurysm had not been seen previously in this setting.

References:

Dziewas R, Konrad C, Drager B, et al: Cervical artery dissection - clinical features, risk factors, therapy and outcome in 126 patients. J Neurol 2003; 250:1179-1184.

Komiyama M, Ishiguro T, Matsusaka Y, Yasui T: Simultaneous dissection of intra- and extracranial vertebral artery. Report of two cases and review of literature. Acta Neurochir 2002; 144:729-733.

Nadgir RN, Loevner LA, Ahmed T, et al: Simultaneous bilateral internal carotid and vertebral artery dissection following chiropractic manipulation: case report and review of the literature. Neuroradiology 2003; 45:311-314.

Nishioka T, Maeda Y, Tomogane Y, Nakano A, Arita N: Unexpected delayed rupture of the vertebral - posterior inferior cerebellar artery aneurysms following closed head injury. Acta Neurochir 2002; 144:839-845.